

## Research Article

### INTENSITY OF FORAGING ACTIVITY IN SUBTERRANEAN TERMITES AT MULAKALAPALLY FOREST REGION, KHAMMAM DISTRICT, ANDHRAPRADESH INDIA

Nageswara Rao. A\* and Ch. Sammaiah

Environmental Biology Laboratory, Department of Zoology, Kakatiya University, Warangal - 506 009, INDIA

\*Corresponding author E-mail: [drapkanagesh@gmail.com](mailto:drapkanagesh@gmail.com)

#### ABSTRACT

Seasonal foraging activity of subterranean termites on forest plants were observed Mulakalapally forest region, Khammam (A.P). Two distinct foraging patterns were identified in the species of termites. *Microtermes Sp* and *Coptotermes Sp* have strong foraging activity high peak in wet season. Whereas *Odontotermes Sp* and *Macrotermes Sp* have strong foraging activity peak in the dry season. The foraging activity depends upon Environmental factors. The foraging activity was observed in the galleries at the temperature to 17°-25°C.

#### KEY WORDS

Mulakalapally forest region, Temperature, *Odontotermes obesus* foraging activity

#### STUDY AREA:

The present investigation was carried out in Mulakalapally forest region, Khammam district in Andhra Pradesh, India; It lies in between latitude 17°36' and 18°38', North and 80°21' East of Greenwich longitudes. The annual average temperature was recorded 25°C and annual precipitation was 990mm. Soil are typically acidic (pH 4.0-7.5) the soil are high in organic matter.

#### INTRODUCTION:

Termites play a key role in tropical and subtropical forest ecosystems as decomposers of the largest part of dead organic matter, playing a significant role in soil formation. At the same time, termites are dangerous pests to agricultural crops and forestry. Studies of termites ecology have been carried out mainly in rain forests (Abe 1978, Abe Matsumoto 1979, Eggleton et al., 1999) .In semi deciduous forests the population has been studied (Sing, Singh 1981, Inoue et al., 2006,

Anichkin et al., 2007, Vu et al 2007). Termites in forest ecosystem of national Park in Vietnam, Belyaeva, A.V. Tiunov (2010). Termites feed on plants in almost every form living, freshly dead in various stages of decomposition and humus. Their feeding habits are reviewed in Adamson (1943) (Lee & Wood 1971).The pattern of damage varies with species, climate and area, it is widely assumed that rarely damage indigenous trees (Harris, 1955; Lee& Wood 1971; Tho 1992).

Termite damage to living trees fall under two distinct categories mainly the damage by monophagous termites which have restricted distribution and colonies are confined to single species of trees, and the damage by polyphagous termites that the forage over a wide range of trees (Das 1958; Roonwal, 1979; Sammaiah, 2002; Sammaiah and Reddy, 2005).

Termites also know one of the prominent ecosystem engineers; (Jones et al 1994, Pardesh, M and A.K. Prusty 2010). Which modify the soil properties by constructing huge mounds and long

subterranean galleries and which provide animals and plants, with heterogeneous habitats. Termites as ecosystem engineers, they contribute significantly to ecosystem processes including carbon fluxes, soil formation and soil conditioning (Lee and Wood 1971, Bignell et al 1997, Bignell and Eggleton 2000).

**MATERIALS AND METHODS**

**Survey and Sampling**

The study was carried out in the deciduous forest region of Mulakalapally in Khammam district A.P. I was surveyed during June, 2008 to May 2010 for termite attack to trees, decomposition of litter foraging activity of termites. From the general survey it was clear that the general activity of termites was more during mornings and late evenings. Hence, the studies were mainly carried out during that time. Trees susceptibility to termite damage was checked. Samples of unhealthy looking trees from selected forest area were sampled. Termites were collected from different parts of the trees. Furthermore, effects were also made to sample all possible habitats. Intensive observations were made on ground, trees, branches, twigs, dead stumps, woody litter and leaf litter for presence of termites. Tunnels, foraging triats on ground, trees or under leaves and debris were also searched. Termites were collected by forceps information in relevant to the specimen, were recorded in writing and also photographed, specimen were preserved in 80% alcohol. The trees were closely and carefully

inspected for signs of termite foraging activity such as earthen sheet covering, run-ways and galleries on living tree-trunks, dead trees and roots were carefully recorded.

The infestation of termites were recorded based on the penetration in to wood and covering of earthen sheet following the slightly modified method of Esenther and Beal (1974) and Beal et al. (1979) which is as follows

- 0 = No infestation
- +=Plastering of earthen sheet of about 5% of the stem of the trees
- ++=Plastering of earthen sheet of about 25% of the stem of the trees
- +++ =Plastering of earthen sheet of about 35% of the stem of the trees
- ++++=Plastering of earthen sheet of about 65% of the stem of the trees
- +++++=80% portion of the trees was covered with earthen sheet

**Photometric method:** In Mulakalapally forest region, a large amount of forest litter falls regularly to the forest floor. In the present study the rate of disappearance of leaf litter due to termite feeding was estimated by visual and photometric methods has been described by Heath et al., 1964, but without using the litter bags.

**Table: 1 Termites infestation to tress in Mulakalapally forest region**

Termite Species	Plant species	Intensity of damage	Damage of Plant Parts
<i>Odontotermes brunneus</i> (Hagen)	<i>Polyalthia longifolia</i>	+++++	stem (live )
	<i>Tectona grandis</i>	++++	dead stem
	<i>Terminalia arjuna</i>	++++	dry log
	<i>Miliusa tomentosa</i>	++++	stem (live)
	<i>Albizia amara</i>	+++	dry wood
	<i>Dalbergia paniculata</i>	+++	dry wood

	<i>Hardwickia binata</i>	+++	stem (live)
	<i>Madhuca indica</i>	+++	dead wood
	<i>Buchanania lanzan</i>	++++	stem (live)
	<i>Acacia chundra</i>	+++	dry part of tree
	<i>Largerstroemia parviflora</i>	+++++	dry part of branches
<i>Odontotermes obesus</i> (Rambur)	<i>Bridelia retusa</i>	+++++	dead wood
	<i>Ixora pavetta</i>	++++	live bark
	<i>Chloroxylon swietenia</i>	++++	stem (dry)
	<i>Senna auriculata</i>	+++	dry wood
	<i>Madhuca indica</i>	+++++	dry branches of tree
	<i>Albizia odoratissima</i>	++++	stem (dry tree)
	<i>Azadirachta indica</i>	+++	stem (live)
	<i>Lannea coromandelica</i>	++++	dead wood
	<i>Alangium salvifolium</i>	+++	dead wood
	<i>Schleichera oleosa</i>	+++	dead stem
	<i>Grewia tilaefolia</i>	++++	fallen leaves
	<i>Morinda pubescens</i>	+++	stem (live)
	<i>Anacardium occidentale</i>	+++++	stem (live)
	<i>Anogeissus latifolia</i>	++++	dry bark
	<i>Senna auriculata</i>	++++	stem (live)
	<i>Milusa tomentosa</i>	++++	stem (dead)
	<i>Tectona grandis</i>	+++++	dry stem
<i>Odonotermes indicus</i> (Thakur)	<i>Albizia odoratissima</i>	+++	dead wood
	<i>Milusa tomentosa</i>	+++	dead wood
<i>O. feae</i> (Wasmann)	<i>Anogeissus latifolia</i>	++++	stem (dead)
	<i>Lagerstroemia parviflora</i>	+++++	dry wood
	<i>Senna auriculata</i>	+++	bark (live)
	<i>Schleichera oleosa</i>	+++++	stem (live)
	<i>Syzygium cumini</i>	++++	stem (live)

	<i>Lagerstroemia parviflora</i>	+++++	dead wood
	<i>Acorus calamus</i>	++++	stem (live)
	<i>Buchanania lanzan</i>	++++	stem (live)
	<i>Terminalia alata</i>	++++	dead wood
<i>O.wallonensis</i> (Wasmann)	<i>Chloroxylon swietenia</i>	++	stem (live)
	<i>Boswellia serrata</i>	++	stem (live)
	<i>Milusa tomentosa</i>	+++	bark (live)
	<i>Anogeissus latifolia</i>	++++	stem (live)
	<i>Madhuca indica</i>	+++	stem (live)
	<i>Tectona gradis</i>	+++	stem (live)
<i>O.redmanni</i> (Wasmann)	<i>Cleistanthus collinus</i>	+++	dead wood
	<i>Premna tomentosa</i>	+++	dead (wood)
	<i>Buchanania lanzan</i>	+++	dead stem
	<i>Anogeissus latifolia</i>	++++	stem (live)
	<i>Syzygium cumini</i>	+++	dead (stem)
	<i>Tectona grandis</i>	+++	stem (live)
	<i>Polyalthia longifolia</i>	++++	dead wood
<i>O. guptai</i> (Roonwal and Bose)	<i>Wrightia tinctoria</i>	+++	stem (dead)
	<i>Senna auriculata</i>	+++++	stem (live)
	<i>Terminalia alata</i>	+++++	stem (live)
	<i>Tectona grandis</i>	+++	stem (dead)
<i>Microtermes obesi</i> (Holmgren)	<i>Terminalia bellerica</i>	+++	dead stem
	<i>Manilkara hexandra</i>	+++	dead (wood)
	<i>Cassia fistula</i>	+++	stem (dead)
	<i>Premna tomentosa</i>	++	dry bark
	<i>Limonia acidissima</i>	+++	dead (stem)
<i>Microcerotermes beesoni</i> (Snyder)	<i>Terminalia alata</i>	+++	live stem
	<i>Calycapteris floribunda</i>	+++++	dead wood
<i>O. boveni</i> (Thakur)	<i>Polyalthia longifolia</i>	++++	dry twig

<i>Heterotermes indicola</i> (Wasmann)	<i>Anogeissus latifolia</i>	+++	dead wood
	<i>Dalbergia sissoo</i>	+++	dead wood
<i>Coptotermes hemi</i> (Wasmann)	<i>Lagerstroemia parviflora</i>	++++	stem (dead)
	<i>Tectona grandis</i>	++++	root and stem (live)
	<i>Cleistanthus collinus</i>	++	stem (live)
	<i>Lannea coromandelica</i>	++	stem (live & dead)

## RESULTS AND DISCUSSION:

Various termite species were collected from different trees, mounds, the fallen tree pieces of logs, dead tree stumps, twigs on the forest floor in the mulakalapally forest area. They are belonging to two families viz., Termitidae and Rhinotermitidae. In Termitidae *Odontotermes boveni* (Thakur), *Odontotermes brunneus* (Hagen), *Odontotermes feae* (Wasmann), *Odontotermes guptai* (Roonwal and Bose), *Odontotermes indicus* (Thakur), *Odontotermes obesus* (Rambar), *Odontotermes redemanni* (Wasmann), *Odontotermes wallonensis* (Wasmann), *Macrotermes convulsionaries* (Konig), *Microceretermes beelsoni* (snyder) and *Microtermes obesi* (Wasmann), and in Rhinotermitidae, *Coptotermes hemi* (Wasmann), and *Heterotermes indicola* (Wasmann).

The nature of damage of different species of termites in the different forest trees. The damage of *Odontotermes* species was recorded in the form of nibbling on dead as well as on live bark of both stem and root underneath the cover of earthen sheet and runways. The attack of *Odonototermes* species occurred usually at the basal part of the trunk. The infestation although not severe, was localized resulting in the formation of irregular cavities or grooves of various sizes on the surface of the trunk. The earthen sheet of *O. brunneus* on *Milius tomentosa* was recorded up to 6 meters height. Its damage was recorded on dead part of stem. It is very intensively damage some more plants were *Terminalia arjuna*, *Albiziaamara*, *Milliusa tomentos*. *Dalbergia paniculats*, *Hardwickia binata*, *Buchanania lanzan*, *Madhuca indica*, *Acacia chundra*, *Lanorgerstromia parviffora*.


The intensity of different termites damage to different trees in Mulakalapally forest region is presented ( table -1) The damage of *O. brunneus* was recorded maximum 80% and minimum 35%.The following trees were infested by this species viz..

*Tectona grandis*, *Largerstroemia parviflora*, *Termanalia arjuna*, *Miliusa tomantossa*, *Buchanania lanzan*. *Albiziamara*, *Dalbergia paniculata*, *Harwickia binata*, *madhuca indiacca* and *Accacia chundra*. The damage of *O. obesus* was recorded maximum 80% and minimum 35%.Its damage was recorded on trees *Brideliavetura*, *Madhuka indiacca*, *Anacardium occidentalis*, *Tectona grandius*, *Ixorapavatta*, *Cloroxylon swietanliq*. *Albizia odoratisima*, *Lannea coromandelica*, *Grewiatilac folia*, *Anogeissus latifolia*, *Miliusa tomentosa* and *Acassia auriculata*. *Azadaracta indiacca*, *alangium salvifolium*, *Schleichera oleora*, *Morinda pubescene*, *lagerstroemia parviflora*, *Schleichera oleora*.

The damage of *O. guptai* was recorded maximum 80% and minimum 35%.Its damage was recorded on trees *Cassia auriculata*, *Terminalia tomentosa*, *Wrightinetoria*, *Terminalia arjuna* and *Tectona grandis* The damage of *ofeae* was recorded maximum 80% and minimum 35%.The following trees were infested by this species viz. *Anogeissus latifolia*, *Syrigium cuminilive*, *Acoras calamus*, *Terminalia tomentosa* and *buchananai lanzan*, *Cassia auriculata*. The damage of *O. wallonensis* was recorded maximum 65% and minimum 35%. Its damage was recorded on trees *Buchanania lazan*, *Anogeissus latifolia*, *Boswellia serrata*, *tectona grandis*, *Chloroxylon swietenia* *Madhuca*

*indica Miliusa tomentosa* The damage of *O. indicus* was recorded maximum 35% and Minimum 35%.Its damage was recorded on trees *Miliusa tomentosa*, *Aleberzia odoratissima* The damage of *O. redmanni* was recorded maximum 65% and minimum 35%. The following trees were infested by this species viz. *Szygium cumini*, *Premna tomentosa*, *Anogeissus latifolia*, *Buchanania lanzan*, *Cleistanthus collinus*, *Tectona grandis*.The damage of *O. boveni* was recorded maximum 65% and minimum 35%.The following trees was infested by this species viz. *Lanorgerstromia parviflora*. *Anogeissus latifolia*. The damage of *Microtermes obesi* was recorded maximum 80% and minimum


35%.Its damage was recorded on *Limonia acidissima*, *Manilkara hexandra*, *Cassia fistula*, and *Premma tomentosa* and *Terminalia bellrica* the damage of *Microtermes beesoni* was recorded maximum 80% and minimum 35%. The following trees were infested by this species viz. *Terminalia tomentosa*, *Calycapteris floribunda*. The damage of *Heterotermes indicola* was recorded maximum 80% and minimum 35%.Its damage was recorded on *Anogeissus latifolia*. The damage of *Captotermes hemi* was recorded maximum 65% and minimum 25%.Its damage was recorded on *Lagerstromia parviflora*, *Tectona grandis*, *Cleistanthus collinus*, and *Lanne coromandelica*.



The percentage of damage caused by *Odontomermes* sp in Mulakalapally forest region is presented in (Fig.1). Maximum infestation was recorded in month of April 2010 and minimum infestation was recorded month February 2010. The infestation in June 2008 (14.15%), it was slightly increased in month of July and slightly decreased in month of August to October 2008. Again it slightly increased in month of November. Again decreased in month of December 2008 then

it was slightly increased from January to February 2009. Again it decreased in March 2009 again suddenly increased in April 2009. Then it gradually decreased from May to September 2009, and slightly increased October and decreased November 2009. Again it was increased December 2009. It was gradually increased from January 2010 to march 2010, and slightly increased in month of April 2010 and again decreased in May 2010.





Overall percentage of damage caused by different species of termites in Mulakalapally forest region. The damage caused by different species in Mulakalapally forest region is presented in (Fig.2). The damage caused by *O. brunneus* (16%) followed by *O. obesus* (24%) *O. redemanni*, (13%) *O. wallonensis* (12%) *O. guptai* (5%) *O. feae* (2%)

*O.boveni* (2%) *Microtermes obesi* (3%) *Heterotermes indicola* (6%) *Coptotermes hemi* (4%), *Microcerotermes beelsoni* (6%) The results indicate that maximum damage caused by *O. obesus* and minimum damage caused by *O. boveni*, *O. feae* and *Microtermes obesi* in Bhadrachalam forest region

## CONCLUSION

Termites are widely distributed in Mulakalapally forest region. They play in an important role in decomposition procession in forest and few termites are causing damage to the plants.

The total 22.6% of forest trees damaged by termites in Mulakalapally forest, our results supported by Harris, 1955 and Lee & Wood, 1971. Termites rarely cause significant damage on tree species.

Our results may shed same light on some of the mechanisms responsible for tree resources exploitation by termites. With our results conclude that presence of termite galleries or earthen sheet on trees could be determined more by resources availability than resource quality. Termites selecting large trees merely because they are easier to be encountered during random foraging.

Results indicated that the foraging activities of subterranean termites in rainy season and winter season was the highest slightly decreased at the end of rainy season. Climatic factors especially rainfall declined soil temperature and increase soil moisture and relative humidity might influenced the foraging patterns and increasing foraging activity of subterranean termites these results supported by Lafage et al ., 1976; Black and wood, 1989.

Comparing the foraging activity throughout the year between different termite species found in the forest revealed that *Odontotermes obesus* showed the highest in foraging activity followed by *O. brunneus*, *O. redemanni*, *O. wallonensis*, *H. indicola*, *C. hemi*, *O. feae* and *O. guptai*.

## ACKNOWLEDGEMENT

The authors are grateful to Dr. M. Ahmed, Head, Division of Forest Entomology, Forest Research Institute, Dehradun for identification of termite species.

## REFERENCES:

- Esenther, G.R and Beal, R.H (1974). Attractant mirex betf suppresses activity of *Reticulitermes spp* journ. Eco. Entomol. 67(1): 85-88..
- Beal, R.L. Bultman, J.D. Amburgey, T.L. and Roberts, D.R. (1979). Resistance of wood from pare quaff treated southern pines to subterranean termites Decay Fungi and Marine Borers, Forest products journal, 29(4): 35-38.
- Anichkin ,A.E., Belyaeva,N.V Dogobrod , I.G., et al., soil Microarthropods and Macro fauna 2007, in monsoon tropical forests of CatTien and Bi dup-Nui Ba National Parks,Southern veitnam Izv. Akad Nauk Ser .Bio.,2007 vol.34,no5,pp .598-607.
- Harris, W.V. 1971. Termites their recognition and control. 2<sup>nd</sup> editin long man, Harlog. activity of soil animals in the breakdown of leaves pedobiologia 4, 80-87.
- Sammaiah, Ch. and Reddy, M.V. (2005). Assessment of biodiversity of termites and their damaged trees in Semi – arid tropical urban systems. Eco. Env. & Cons., 11(2): 87 -96.
- Sammaiah, Ch. (2002). Termite attacking fruit crops in emi and tropical agroecosystem. Modern trends in Entomological Research in India, PP. 73-76.
- Matsumoto,T. and Abe, T. (1979). The role of termites in a equatorial rain forest ecosystem of west Malaysia. II. The role of termites in the decomposition of leaf litter on the forest floor. Ecologies 38: 261 – 274.
- Inoue,T., Takematsu , Y., Yamada,A., et al ., Diversity and abundance of termites along an altitudinal gradient in Khao Kitchagoot National park , Thailand ,j.Trop.ecol.2006 vol.22,pp.609-612.
- Harris W.V. (1955). Termites and Forestry. Emp. For. Rev. 34 : 160 – 166
- Das, G.M. (1958). Observation on the termites effecting tea in north – East Indian and their control. Indian J. Agric, Science. Q28 (4): 553-560.
- Eggleton, p., Homathevi, R ., Jones,D.T.,et al., Termites assemblages , forest Disturbance and Greenhouse Gas Fluxes in Sabh ,East Malaysia, phi.trans.R Soc .L.B. 1999, vol 354,PP.1791-1802.
- Tho, Y.P. (1992). Termites of peninsular Malaysia (Kirton) L.G.Ed. Malayan Forest records, No. 36: 224pp forest research institute. Malaysia Kepong.
- Lee & Wood 1971. Adamson, A.M. (1943) .Termites and the fertility of soils, *Trop. Agr.* 20(6): 107-112.
- Roonwal, M.L (1979). Termite life and termite control in south Asia. IX : PP. 1-77.
- Lee K.e, Wood TG (1971). Termites and soils. London, Academic Press P.251 New york
- Belyaeva and A.V. Tiunov (2010) Termites (Isoptera) in forest ecosystem of cattien National Park (Southern Vietnam) Journal Biology Bulletin 37, No. 4 pp. 374-381.
- Singh, U.R. and Singh, J.S. (1981). Population structure and mound architecture of the termites of a tropical deciduous forest of Varanasi, India. Pedobiologia 22 : 213 – 223.
- Vu,Q.M.,Nguyen,H.H., and Smith R.L, (2007) The termites (Isoptera) of Xuan Son National park,Northern Vietnam Pan- Pacific Entomol ., vol.83,pp.85-94.
- Pardesh:, M. and B.A.K. Prusty, 2010. Termites as ecosystem engineers and potentials for soil restoration curr. Sci., gg: 11-11.
- Abe, T and T. Matsumoto. (1978). The role of termites in an equatorial rain forest ecosystem of west Malaysia II leaf litter consumption on the forest floor. *Oecologia* 38: 261-274
- Lafage, J.P., M.I. Haverty and W.L. Nutting (1976). Environmental factors co related with the foraging behavior of desert subterranean termite, *Gnathami termes perplexus* (Banks). (Isoptera : Termitidae). Sociobiology, 2: 155-169.
- Jones, C.G., Lawton, J.H. and shachak, M. (1994). Organisms as ecosystem engineers. *Oikos* 69(3): 373 – 386. La – Fage, J.P. and Nutting, W.L. 1978. Nutrient dynamics of termites. Production Ecology of Ants and Termites (Brain, M.V., ed). Cambridge University Press, Cambridge. PP. 165 – 232.
- Black, H.I and T.G. Wood (1989). Effect of cultivation on vertical distribution of *Macrotermes spp.* (Isoptera Termitidae : Macro termitinae) is soil at Mokwa, Nigeria, Sociobiology 15, 133-138.
- Black, H.I and T.G. Wood (1989). Effect of cultivation on vertical distribution of *Macrotermes spp.* (Isoptera Termitidae : Macro termitinae) is soil at Mokwa,Nigeria, Sociobiology 15, 133-138.
- Bignell DE, Eggleton P (2000). Termites in ecosystems. In Abe, T., Bignell, D.E. & Higashi, M. (Eds) Termites, Evolution, Sociality, Symbioses, Ecolozy, Dordrecht, the Netherlands, Klower Acadimic Publishers, PP. 363 – 388.
- Bignell, D.E., Eggleton, P., Nunes, L. & Thomas, K.L. 1997 Termites as mediators of Carbon fluxes in tropical forest: budgets for carbon dioxide and Methane emissims. In: Watt, A.D., Stork, N.E &



- Hunter, M.D. (eds) Forest and Insects Chapman and Hall, London: 109 - 134.
- Roonwal, M.L. (1970). Termites of the oriental region in K. Krishna and F.M. Weesner, eds. Biology of termites. Academic Press, New York, PP. 320-391.
  - Das, G.M. (1958). Observation on the termites effecting tea in north - East Indian and their control. Indian J. Agric, Science. Q28 (4): 553-560.